



## *TopoCellGen*

Generating Histopathology Cell Topology  
with a Diffusion Model




### Oral Presentation


Meilong Xu, Saumya Gupta, Xiaoling Hu, Chen Li, Shahira Abousamra,  
Dimitris Samaras, Prateek Prasanna, **Chao Chen**

# What is Cell Topology?


Histopathology Image




Intra-class Distribution



Histopathology Image



Inter-class Relationship



Tumor/Epithelial


Lymphocyte

Stromal Cell

# Why Cell Topology?

Diagnosis depends on who a cell's neighbors are.

Single-class cell distribution




Normal glandular structure

Grade 4 of Gleason

Wikimedia Commons

Multi-class cell relationships




Mlecnik et al. *Immunity*, 2016<sup>1</sup>

# Cell Topology Generation


$$c = [c_0, c_1, c_2]$$

- Lymphocyte
- Tumor/Epithelial
- Stromal Cell

Cell Count  
Per Class




- Cell count and density
- Intra-class cell distribution
- Inter-class cell interactions




Aiming to generate accurate  
cell topology!

# Method: Overview



# Method: Ensure accurate count per class

- Pixel-level enforcement
- Maintain the count of cells for each cell type




$$\mathcal{L}_{count} = \frac{1}{|n|} \sum_{i=1}^n \left| \frac{\sum b(\hat{x}_0^t)^{(i)}}{\delta} - \frac{\sum b(x_0^t)^{(i)}}{\delta} \right|$$


# Topological Feature

- **Cell topology** – Here mainly represents the holes
- Use **Persistent Homology**<sup>2</sup> to capture the holes.


H&E Patch




Cell Layout



*Increasing Disk Radius  $r$*




Represent in a Persistence Diagram (Dgm):




# Matching Structures: Holes

- Target: Ref. and Syn. have **similar intra/inter-class topological patterns**.
- Find the **correspondence** between the Ref. and Syn. structures.
- The matching is based on the **Wasserstein distance<sup>3</sup>**.




# Method: $\mathcal{L}_{intra}$

- Matched structures should have **similar spatial patterns of same type of cells**.
- e.g. have similar values at the matched locations in **class-wise distance transform map**.




# Method: $\mathcal{L}_{inter}$

- Matched structures should have **similar spatial patterns** of different types of cells.
- e.g. have similar values at the matched locations in a **combined** distance transform map.




# New Metric: Topological Fréchet Distance

- Conventional FID measures feature similarity, but fails to capture complex spatial and topological cell distributions.




# Qualitative Results


H&E  
Image



Reference  
Cell Layout



Synthetic  
Cell Layout



# Quantitative Evaluation: Generation Quality

- Conduct experiments on BRCA-M2C and Lizard datasets.

	Method	FID ↓	Lym. ↓	Epi. ↓	Stro. ↓	Neu. ↓	Pla. ↓	Eos. ↓	Con. ↓	TCE ↓	TopoFD ↓	MMD ↓
BRCA-M2C	ADM [11]	1.150	13.757	40.230	15.491	–	–	–	–	22.465	133.012	0.732
	TMCCG [2]	0.634	11.503	34.032	12.907	–	–	–	–	19.687	89.252	0.635
	Spatial Diffusion [44]	0.263	10.852	35.954	13.496	–	–	–	–	20.806	97.584	0.589
	<i>TopoCellGen</i>	<b>0.005</b>	<b>2.090</b>	<b>3.824</b>	<b>2.468</b>	–	–	–	–	<b>5.192</b>	<b>69.354</b>	<b>0.421</b>
Lizard	ADM [11]	0.059	16.508	11.796	–	1.123	4.328	1.598	10.737	23.964	65.910	0.783
	TMCCG [2]	1.093	15.548	10.011	–	2.376	4.293	1.872	11.643	22.604	63.120	0.667
	Spatial Diffusion [44]	0.137	10.740	9.062	–	3.040	6.552	2.173	11.225	20.606	79.591	0.883
	<i>TopoCellGen</i>	<b>0.027</b>	<b>6.155</b>	<b>6.560</b>	–	<b>1.022</b>	<b>2.982</b>	<b>1.167</b>	<b>7.288</b>	<b>11.590</b>	<b>31.607</b>	<b>0.536</b>

# Quantitative Evaluation: Downstream Performance

- Downstream Task: cell detection and classification

Data	Method	F1-Score ↑				
		Lymphocytes	Epithelial	Stromal	Mean	Detection
Real.	UNet	0.569 ± 0.010	0.736 ± 0.012	0.507 ± 0.015	0.604 ± 0.011	0.857 ± 0.006
Real+Syn. (Rand)		0.549 ± 0.009	0.693 ± 0.014	0.472 ± 0.016	0.571 ± 0.013	0.848 ± 0.008
Real+Syn (TMCCG)		0.650 ± 0.007	0.768 ± 0.010	0.511 ± 0.012	0.643 ± 0.009	0.852 ± 0.005
Real+Syn (SpaDM)		0.647 ± 0.006	0.797 ± 0.003	0.554 ± 0.011	0.666 ± 0.007	0.853 ± 0.005
Real+Syn ( <i>TopoCellGen</i> )		<b>0.656 ± 0.003</b>	<b>0.803 ± 0.005</b>	<b>0.574 ± 0.004</b>	<b>0.678 ± 0.004</b>	<b>0.860 ± 0.004</b>
Real.	MCSpaNet	0.615 ± 0.008	0.777 ± 0.010	0.540 ± 0.013	0.644 ± 0.009	0.855 ± 0.005
Real+Syn. (Rand)		0.578 ± 0.009	0.756 ± 0.012	0.502 ± 0.014	0.612 ± 0.010	0.851 ± 0.006
Real+Syn (TMCCG)		<b>0.678 ± 0.006</b>	0.800 ± 0.005	0.522 ± 0.014	0.667 ± 0.007	0.853 ± 0.004
Real+Syn (SpaDM)		0.639 ± 0.005	0.804 ± 0.007	0.563 ± 0.012	0.669 ± 0.006	0.855 ± 0.005
Real+Syn ( <i>TopoCellGen</i> )		0.652 ± 0.004	<b>0.817 ± 0.006</b>	<b>0.582 ± 0.005</b>	<b>0.684 ± 0.004</b>	<b>0.862 ± 0.004</b>

# Conclusion

- The first diffusion-based framework for cell topology generation.
- Introduce a novel topology-aware metric to quantify topological fidelity of synthetic layouts.
- Enhance downstream task performance greatly using augmented image-layout pairs.

# Thank you!

